

Package leaflet: Information for the user
CIALIS 20 mg film-coated tablets
tadalafil

Read the entire leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- Do you have any questions? Contact your doctor or pharmacist.
- Do not pass this medicine on to others, as it has been prescribed only for you. It may be harmful to others, even if their symptoms are the same as yours.
- Do you experience any side effects listed in section 4? Or do you experience a side effect that is not listed in this leaflet? Then contact your doctor or pharmacist.

Contents of this leaflet

1. What is CIALIS and what is it used for?
2. When should you not use this medicine or be extra careful?
3. How to use this medicine?
4. Possible side effects
5. How to store this medicine?
6. Contents of the pack and other information

1. What is CIALIS and what is it used for?

CIALIS is intended for the treatment of adult men with erectile dysfunction. This is due to the order when a man cannot get an erection or cannot maintain an erection long enough that is suitable for sexual activity. CIALIS has been shown to significantly improve the ability to achieve an erection suitable for sexual activity.

CIALIS contains the active substance tadalafil, which belongs to a group of medicines called phosphodiesterase type 5 inhibitors. After sexual stimulation, CIALIS helps relax the blood vessels in the penis, allowing blood to flow into your penis. This results in an improvement of your erection. CIALIS does not work if you do not have erectile dysfunction.

It is important to know that CIALIS does not work without sexual stimulation. You and your partner need to know that foreplay is necessary, just as you would if you were not using a medicine for erectile dysfunction.

2. When should you not use this medicine or be extra careful with it?

When should you not use this medicine?

- You are allergic to tadalafil or any of the other ingredients in this medicine. These ingredients can be found in section 6.

- You are using an organic nitrate in any form or a medicine that releases nitric oxide (such as amyl nitrite). This is a group of medicines (nitrates) used for the treatment of angina pectoris (chest pain). It has been shown that CIALIS enhances the effect of these medicines. If you are using nitrate in any form or are unsure whether you are using it, tell your doctor.
- U have a severe heart condition or have had a heart attack in the last 90 days.
- U have had a stroke in the last 6 months.
- U have low blood pressure or high blood pressure that is not controlled.
- U have ever had loss of vision due to non-arteritic ischemic anterior optic neuropathy (NAION), a condition described as “eye paralysis”. U
- are using riociguat. This medicine is used to treat pulmonary arterial hypertension (this is high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (this is high blood pressure in the lungs due to blood clots). PDE5 inhibitors, such as CIALIS, have been shown to increase the blood pressure-lowering effect of this medicine. If you are using riociguat or are unsure, contact your doctor. When should you be extra careful with this medicine?

Contact your doctor before using this medicine.

Be aware that sexual activity can be risky for patients with a heart condition

Be aware that sexual activity can be risky for patients with a heart condition because this activity requires extra effort from your heart. If you have a heart condition, tell your doctor.

Tell your doctor before taking the tablets if you have any of the following conditions:

- sickle cell anemia (a disorder of the red blood cells),
- multiple myeloma (bone marrow cancer),
- leukemia (blood cell cancer),
- a deformity of your penis,
- a severe liver problem,
- a severe kidney problem.

It is not known if CIALIS is effective in patients who:

- have undergone pelvic surgery, total or partial removal of the prostate gland, where the nerve pathways of the
- prostate have been severed (radical non-nerve-sparing prostatectomy), have undergone.

If you suddenly notice a decrease or loss of vision, or if your vision is distorted or blurred while using CIALIS, stop using CIALIS and contact your doctor immediately.

Sudden onset of decreased hearing or hearing loss has been observed in some patients after using tadalafil. Although it is not known if these cases are directly due to tadalafil, you should

contact your doctor as soon as you experience sudden onset of decreased hearing or hearing loss notes.

CIALIS is not intended for use by women.

Children and adolescents under 18 years

CIALIS is not intended for use by children and adolescents under 18 years.

Are you taking any other medicines?

Are you taking any other medicines besides CIALIS, have you recently done so, or is there a possibility that you will soon take other medicines? Tell your doctor.

Do not take CIALIS if you are already taking nitrates.

Some medicines are affected by CIALIS or can affect how CIALIS works. Tell your doctor or pharmacist if you are already taking the following medicines:

- an alpha-blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia),
- other medicines to treat high blood pressure,
- riociguat,
- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia), medicines such as ketoconazole tablets (used to treat a fungal infection) and protease inhibitors for the treatment of AIDS or HIV infection,
- phenobarbital, phenytoin, and carbamazepine (medicines used to treat seizures),
- rifampicin, erythromycin, clarithromycin, or itraconazole,
- other treatments for erectile dysfunction.

What should you be aware of with drinking and alcohol?

See section 3 for information on the effect of alcohol use. Grapefruit juice may affect how well CIALIS will work and should be taken with caution. Ask your doctor for more information about this.

Fertility

In treated dogs, there was reduced sperm production in the testes. A reduction in sperm has also been observed in some men. It is unlikely that these effects will lead to a lack of fertility.

Driving and using machines

Some men who took CIALIS in clinical studies reported dizziness. Carefully check how you react to the tablets before driving or operating machinery.

CIALIS contains lactose

If your doctor has told you that you have an intolerance to some sugars, contact your doctor before taking this medicine.

CIALIS contains sodium

This medicine contains less than 1 mmol (23 mg) of sodium per tablet, that is to say, it is essentially 'sodium-free'.

3. How to use this medicine?

Always use this medicine exactly as your doctor has told you. Are you unsure about the correct use? Then contact your doctor or pharmacist.

CIALIS tablets are intended for oral use by men only. Swallow the tablet whole with some water. You can take CIALIS with or without food.

The recommended starting dose is one 10 mg tablet before sexual activity. However, you have been given the dose of one 20 mg tablet because your doctor has decided that the recommended dose of 10 mg is too weak for you.

You can take a CIALIS tablet at least 30 minutes before sexual activity. CIALIS can still be effective up to 36 hours after taking the tablet.

You must not use CIALIS more than once a day. CIALIS 10 mg and 20 mg are intended for use before anticipated sexual activity and are not recommended for continuous daily use.

It is important to know that CIALIS does not work without sexual stimulation. You and your partner should know that foreplay is necessary, just as you would do if you were not using a medication for erectile dysfunction.

Drinking alcohol can affect your ability to get an erection and can temporarily lower your blood pressure. If you have taken CIALIS or plan to take CIALIS, avoid excessive drinking (blood alcohol level of 0.08% or higher), as this can increase the risk of dizziness when standing up.

Have you used too much of this medicine?

Inform your doctor. You may experience side effects as described in section 4.

Do you have any other questions about the use of this medicine? Then contact your doctor or pharmacist.

4. Possible side effects

Like any medicine, this medicine can have side effects, although not everyone gets them. These side effects are usually mild to moderate in nature.

If you experience any of the following side effects, stop taking CIALIS and seek immediate medical help:

- Allergic reaction including rash (occurs sometimes).
- Chest pain - do not use nitrates but seek immediate medical help (occurs sometimes).
- Priapism, a prolonged and possibly painful erection after taking CIALIS (rarely occurs). If you have such an erection and it lasts longer than 4 hours, contact your doctor immediately.

- U suddenly cannot see anymore (rare), distorted, blurred, hazy central vision or sudden reduced vision (frequency not known).

Other reported side effects include:

Common (occur in less than 1 in 10 patients)

- headache, back pain, muscle pain, pain in arms and legs, facial flushing, nasal congestion and digestive problems.

Sometimes (occur in less than 1 in 100 patients)

- dizziness, abdominal pain, nausea (vomiting), reflux, blurred vision, eye pain, difficulty breathing, presence of blood in urine, prolonged erection, palpitations, rapid heartbeat, high blood pressure, low blood pressure, nosebleeds, tinnitus, swelling of hands, feet or ankles, and feeling tired.

Rare (occur in less than 1 in 1000 patients)

- fainting, seizures and transient memory loss, swelling of the eyes, red eyes, sudden decrease or loss of hearing, hives (itchy red welts on the skin surface), bleeding from the penis, presence of blood in semen, and increased sweating.

A heart attack and stroke have also been rarely reported in men taking CIALIS. Most of those men already had heart problems before taking this medication.

There have been rare reports of partial, temporary, or permanent decrease or loss of vision in one or both eyes.

Some other side effects that rarely occurred in men taking CIALIS and were not seen during clinical trials include:

- migraine, swollen face, severe allergic reaction causing the face and throat to swell, severe skin rash, certain disorders related to the blood supply to the eyes, irregular heartbeat, angina, and sudden cardiac death.
- distorted, blurred, hazy central vision or sudden reduction of vision (frequency not known)

The side effect dizziness is more frequently reported in men over 75 years using CIALIS.

Diarrhea is more frequently reported in men over 65 years using CIALIS.

Reporting side effects

If you experience side effects, contact your doctor or pharmacist. This also applies to possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system as mentioned in Appendix V. By reporting side effects, you can help us obtain more information about the safety of this medicine.

5. How to store this medicine?

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date. You can find it on the box and the blister pack after EXP. It includes a month and a year. The last day of that month is the expiry date.

Store in the original packaging to protect from moisture. Store below 30°C.

Do not flush medicines down the sink or toilet and do not throw them in the trash. Ask your pharmacist what to do with medicines you no longer use. If you dispose of medicines in the right way, they are destroyed responsibly and do not enter the environment. justified.

6. What substances are in this medicine?

The active substance in this medicine is

- tadalafil . Each tablet contains 20 mg tadalafil ..
- Tablet core:

lactose monohydrate (see section 2), croscarmellose sodium , hydroxypropyl cellulose, microcrystalline cellulose, sodium lauryl sulfate sodium lauryl sulfate, magnesium stearate, see section 2 "CIALIS contains lactose".

Film coating: lactose monohydrate, hypromellose, triacetin, titanium dioxide (E171), yellow iron oxide (E172), talc.

What does CIALIS look like and what is in a package?

CIALIS 20 mg is a yellow almond-shaped film-coated tablet and is marked on one side with "C 20".

CIALIS 20 mg is available in blister packs containing 2, 4, 8, 10, or 12 tablets.

Not all pack sizes may be marketed.

Marketing authorization holder and manufacturer

Marketing authorization holder: Eli Lilly Nederland B.V., Papendorpseweg 83, 3528 BJ Utrecht, Netherlands

Manufacturer: Lilly S.A., Avda. de la Industria 30, 28108 Alcobendas, Madrid, Spain

For all information about this medicine, contact the local representative of the marketing authorization holder:

Belgium/Belgique/Belgien

Eli Lilly Benelux S.A./N.V.

Tel/Tel: + 32-(0)2 548 84 84

Bulgaria

TP "Eli Lilly Nederland" B.V.- Bulgaria

tel. + 359 2 491 41 40

Czech Republic

ELI LILLY ČR, Ltd.

Tel: + 420 234 664 111

Denmark

Eli Lilly Denmark A/S

Tel: +45 45 26 60 00

Germany

Lilly Germany GmbH

Tel. + 49-(0) 6172 273 2222

Estonia

Eli Lilly Netherlands B.V.

Tel: +372 6 817 280

Greece

PHARMASERV-LILLY.E.B.E.

Tel: +30 210 629 4600

Spain

Lilly S.A.

Tel: + 34-91 663 50 00

France

Lilly France

Tel: +33-(0) 1 55 49 34 34

Croatia

Eli Lilly Hrvatska d.o.o.

Tel: +385 1 2350 999

Ireland

Eli Lilly and Company (Ireland) Limited

Tel: + 353-(0) 1 661 4377

Iceland

Icepharma hf.

Phone: + 354 540 8000

Italy

Eli Lilly Italia S.p.A.

Tel: + 39- 055 42571

Cyprus

Phadisco Ltd

Tel: +357 22 715000

Latvia

Eli Lilly (Suisse) S.A Representation In Latvia

Tel: +371 67364000

Lithuania

Eli Lilly Lithuania

Tel. +370 (5) 2649600

Luxembourg/Luxemburg

Eli Lilly Benelux S.A./N.V.

Tel/Tel: + 32-(0)2 548 84 84

Hungary

Lilly Hungary Ltd.

Tel: + 36 1 328 5100

Malta

Charles de Giorgio Ltd.

Tel: + 356 25600 500

Netherlands

Eli Lilly Netherlands B.V.

Tel: + 31-(0) 30 60 25 800

Norway

Eli Lilly Norway A.S.

Tel: + 47 22 88 18 00

Austria

Eli Lilly Ltd..

Tel: + 43-(0) 1 711 780

Poland

Eli Lilly Poland Ltd. o.o.

Tel: +48 22 440 33 00

Portugal

Lilly Portugal Products Pharmaceuticals, Lda

Tel: + 351-21-4126600

Romania

Eli Lilly Romania S.R.L.

Tel: + 40 21 4023000

Slovenia

Eli Lilly pharmaceutical company, d.o.o.

Tel: +386 (0)1 580 00 10

Slovak Republic

Eli Lilly Slovakia s.r.o.

Tel: + 421 220 663 111

Finland

Oy Eli Lilly Finland Ab

Puh/Tel: + 358-(0) 9 85 45 250

Sweden

Eli Lilly Sweden AB

Tel: + 46-(0) 8 7378800

United Kingdom (Northern Ireland)

Eli Lilly and Company (Ireland) Limited

Tel: + 353-(0) 1 661 4377

This leaflet was last approved in {MM/YYYY}

More information about this medicine is available on the website of the European Medicines Agency: <http://www.ema.europa.eu>.