

Package leaflet: Information for the user

Levitra 20 mg film-coated tablets
Vardenafil

Read the entire leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- Do you have any questions? Contact your doctor or pharmacist.
- Do not pass this medicine on to others, as it has been prescribed for you only. It may harm them, even if their symptoms are the same as yours.
- Do you experience any side effects listed in section 4? Or do you experience a side effect not listed in this leaflet? Then contact your doctor or pharmacist.

Contents of this leaflet

1. What is Levitra and what is it used for?
2. When should you not use this medicine or be extra careful with it?
3. How to use this medicine?
4. Possible side effects
5. How to store this medicine?
6. Contents of the pack and other information

1. What is Levitra and what is it used for?

Levitra contains vardenafil, a drug from the group of medicines called phosphodiesterase type 5 inhibitors. They are used for the treatment of erectile dysfunction in adult men, a condition that makes it difficult to get or maintain an erection.

At least one in ten men has problems getting or maintaining an erection at some point. There can be physical or psychological causes, or a combination of both. Whatever the cause, due to changes in muscles and blood vessels, there is not enough blood in the penis to make and keep it erect.

Levitra only works if you are sexually stimulated. It reduces the effect of a natural chemical in the body that causes the erection to disappear. Levitra allows an erection to last long enough for satisfactory sexual intercourse.

2. When should you not use this medicine or be extra careful with it?

When should you not use this medicine?

- You are allergic to any of the ingredients in this medicine. You can find these ingredients in section 6. Hypersensitivity symptoms may include rash, itching, swollen face or lips, and shortness of breath.
- You have ever developed a severe skin rash or skin peeling, blistering and/or sores in the mouth after taking Levitra.
- You are taking medicines that contain nitrates, such as glyceryl trinitrate, for the treatment of angina pectoris, or medicines that release nitric oxide, such as amyl nitrite. Taking these medicines together with Levitra can cause your blood pressure to change significantly.

- You are taking ritonavir or indinavir, medicines used to treat human immunodeficiency virus (HIV) infection.
- You are over 75 years old and are taking ketoconazole or itraconazole, medicines for fungal infections.
- You have severe heart or liver problems.
- You are undergoing kidney dialysis.
- You have recently had a stroke or heart attack.
- You have or had low blood pressure.
- Your family has a history of degenerative eye disease (such as retinitis pigmentosa).
- You have ever had a condition where vision loss occurred due to damage to the optic nerve from insufficient blood supply, known as non-arteritic anterior ischemic optic neuropathy (NAION).
- You are taking riociguat. This medicine is used to treat pulmonary arterial hypertension (this is high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (this is high blood pressure in the lungs due to blood clots). PDE5 inhibitors, such as Levitra, have been shown to enhance the blood pressure-lowering effect of this medicine. If you are taking riociguat or are unsure, contact your doctor.

When should you be extra careful with this medicine?

Contact your doctor or pharmacist before using this medicine.

Be extra careful with this medicine

- If you have heart problems. It may be dangerous for you to have sex.
- If you suffer from an irregular heartbeat (arrhythmias) or have a hereditary heart disease that affects the electrocardiogram (ECG).
- If you have a physical condition that affects the shape of the penis. These can be conditions such as angulation, Peyronie's disease, or fibrosis of the corpus cavernosum.
- If you have a condition that can cause erections that do not go away (priapism). These include sickle cell anemia, multiple myeloma, and leukemia.
- When you have a stomach ulcer (also known as a peptic ulcer).
- When you have a bleeding disorder (such as hemophilia).
- If you are using other treatments for erectile dysfunction, including Levitra orodispersible tablets (see section: Are you taking any other medicines?).
- Severe skin reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, have been reported in connection with treatment with vardenafil. Stop using Levitra and seek immediate medical help if you notice any of the symptoms related to these serious skin reactions described in section 4.
- If you suddenly experience a decrease or loss of vision or if your vision is distorted or darkened while taking Levitra, stop taking Levitra and contact your doctor immediately.

Children and adolescents up to 18 years

Levitra is not intended for use by children and adolescents under 18.

Are you taking any other medicines?

Are you taking any other medicines besides Levitra, have you done so recently, or is there a possibility that you will take other medicines in the near future? Then tell your doctor or pharmacist. This also applies to medicines for which you do not need a prescription.

Some medicines can cause problems, particularly:

- nitrates, medicines for angina, or substances that release nitric oxide, such as amyl nitrite ('poppers'). Using these medicines with Levitra can significantly change your blood pressure.
- medicines for the treatment of arrhythmias such as quinidine, procainamide, amiodarone, and sotalol.
- ritonavir or indinavir, medicines for HIV.
- ketoconazole or itraconazole, medicines for fungal infections.
- erythromycin or clarithromycin, macrolide antibiotics.
- alpha-blockers, a type of medication used to treat high blood pressure or an enlarged prostate (such as benign prostatic hyperplasia).
- riociguat.

Do not use Levitra film-coated tablets together with another treatment for erectile dysfunction, including Levitra orodispersible tablets.

What should you pay attention to with food, drink, and alcohol?

- You can take Levitra with or without food but preferably not after a heavy or high-fat meal, as this may delay the onset of action.
- Do not drink grapefruit juice when using Levitra. This can affect the action of the medication.
- Alcoholic beverages can worsen erectile dysfunction.

Pregnancy and breastfeeding

Levitra is not intended for use by women.

Driving and using machines

Levitra can cause dizziness or affect vision in some people. If you feel dizzy or have impaired vision after using Levitra, do not drive or use tools or machines.

3. How do you use this medicine?

Always use this medicine exactly as your doctor or pharmacist has told you. Are you unsure about the correct use? Then contact your doctor or pharmacist.

The usual dosage is 10 mg.

Take a Levitra tablet about 25 to 60 minutes before sexual activity. With sexual stimulation, you may have an erection after about 25 minutes to 4-5 hours after taking Levitra.

- Swallow one tablet with a glass of water.

Do not use Levitra film-coated tablets together with other forms of Levitra.

Do not use Levitra more than once a day.

Inform your doctor if you think Levitra is too strong or not strong enough. He or she may suggest switching to another formulation of Levitra with a different dosage, depending on how well it works for you.

Have you used too much of this medicine?

Men who take too much Levitra may experience more side effects or may develop severe back pain. If you take more Levitra than prescribed, inform your doctor.

Do you have any other questions about the use of this medicine? Then contact your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everyone gets them. Most effects are mild to moderate.

Partial, sudden, temporary, or permanent decrease or loss of vision, or distorted, darkened, or blurred central vision in one or both eyes has been experienced by patients. If this happens, stop using Levitra and contact your doctor immediately. Sudden decrease or loss of hearing has been reported.

Cases of sudden death, rapid or altered heart rhythm, heart attack, chest pain, and problems in the blood circulation in the brain (including temporary reduction of blood supply to parts of the brain and bleeding in the brain) have been reported in men using vardenafil. Most of the men who experienced these side effects had pre-existing heart problems before taking this medicine. It is not possible to determine whether these events were directly related to the use of vardenafil.

The likelihood of experiencing a side effect is described in the following categories:

Very common:

may occur in more than 1 in 10 users

- headache

Common:

may occur in up to 1 in 10 users

- dizziness
- flushing
- stuffy nose or runny nose
- digestive disorder

Sometimes:

may occur in up to 1 in 100 users

swelling of skin and mucous membranes, including swollen face, lips, or throat

- sleep disorder
- numbness and reduced sense of touch
- drowsiness
- effects on vision, red eyes, effects on color vision, eye pain and discomfort, sensitivity to light
- tinnitus, vertigo
- rapid heartbeat or pounding heart
- shortness of breath
- stuffy nose
- heartburn, gastritis, abdominal pain, diarrhea, vomiting, nausea, dry mouth
- increased amount of liver enzymes in the blood

- skin rash, redness of the skin
- back or muscle pain, increased amount of a muscle enzyme in the blood (creatine phosphokinase), stiff muscles
- persistent erections
- malaise

Rarely:

may occur in up to 1 in 1,000 users

- eye inflammation (conjunctivitis)
- allergic reactions
- anxiety
- fainting
- memory loss
- seizures
- increased pressure in the eye (glaucoma), increased tearing
- effects on the heart (such as a heart attack, altered heart rate, or angina)
- high or low blood pressure
- nosebleed
- effect on the results of blood tests for checking liver function
- sensitivity of the skin to sunlight
- painful erections
- chest pain
- temporary reduction of blood supply to parts of the brain

Very rare or not known:

may occur in less than 1 in 10,000 users or the frequency cannot be determined based onis from the available data

- blood in the urine (hematuria)
- bleeding of the penis (penile bleeding)
- presence of blood in the semen (hematospermia)
- sudden death
- bleeding in the brain
- Reddish non-raised, disc-shaped or round spots on the trunk, often with central blistering, skin peeling, sores in the mouth, throat, nose, genitals, and eyes. These severe skin reactions may be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome, toxic epidermal necrolysis)
- Distorted, darkened or blurry central vision or sudden decrease in vision

Reporting side effects

If you experience side effects, contact your doctor or pharmacist. This also applies to possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system as listed in Appendix V. By reporting side effects, you can help us obtain more information about the safety of this medicine.

5. How to store this medicine?

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date. You can find it on the box after 'EXP'. It includes a month and a year. The last day of that month is the expiry date.

Do not flush medicines down the sink or toilet and do not throw them in the trash. Ask your pharmacist what to do with medicines you no longer use. They will be destroyed responsibly and will not enter the environment.

6. Contents of the package and other information

What substances are in this medicine?

- The active substance in this medicine is vardenafil. Each tablet contains 20 mg vardenafil (as hydrochloride).
- The other substances in this medicine are:
Tablet core: crospovidone, magnesium stearate, microcrystalline cellulose, colloidal silicon dioxide (anhydrous).
Film coating: macrogol 400, hypromellose, titanium dioxide (E171), yellow iron oxide (E172), red iron oxide (E172).

What does Levitra look like and how many are in a package?

Levitra 20 mg film-coated tablets are orange with the Bayer cross on one side and the strength (20) on the other side. The tablets are supplied in a blister pack with 2, 4, 8, 12, or 20 tablets. Not all pack sizes may be marketed.

Marketing authorization holder and manufacturer

Marketing authorization holder

Bayer AG
51368 Leverkusen
Germany
Manufacturer

Bayer AG
Kaiser-Wilhelm-Allee
51368 Leverkusen
Germany

For all information regarding this medicine, contact the local representative of the marketing authorization holder:

Belgium/Belgique/Belgien
Bayer SA-NV
Tél/Tel: +32-(0)2-535 63 11

Lithuania
UAB Bayer
Tel: +370 523 36 868

Bulgaria
Bayer Bulgaria EOOD
Tel.: +359-(0)2-424 72 80

Luxembourg
Bayer SA-NV
Tél/Tel: +32-(0)2-535 63 11

Czech Republic
Bayer s.r.o.
Tel: +420 266 101 111

Hungary
Bayer Hungária KFT
Tel.: +36 14 87-4100

Denmark

Malta

Bayer A/S
Tlf: +45 45 23 50 00

Alfred Gera and Sons Ltd.
Tel: +35 621 44 62 05

Germany
Jenapharm GmbH & Co. KG
Tel: +49 (0)3641-87 97 444

Netherlands
Bayer B.V.
Tel: +31-(0)23 799 1000

Estonia
Bayer OÜ
Tel: +372 655 85 65

Norway
Bayer AS
Tlf: +47-23 13 05 00

Greece
Bayer Ελλάς ABEE
Tel: +30 210 61 87 500

Austria
Bayer Austria Ges. m. b. H.
Tel: +43-(0)1-711 46-0

Spain
Bayer Hispania S.L.
Tel: +34-93-495 65 00

Poland
Bayer Sp. z o.o.
Tel.: +48 22 572 35 00

France
Bayer HealthCare
Tél (N° vert) : +33-(0)800 87 54 54

Portugal
Bayer Portugal, Lda.
Tel: +351 21 416 42 00

Croatia
Bayer d.o.o.
Tel: + 385-(0)1-6599 900

Romania
SC Bayer SRL.
Tel: +40-21-529 59 00

Ireland
Bayer Limited
Tel: +353 1 216 3300

Slovenia
Bayer d. o. o.
Tel: +386 (0)1 58 14 400

Iceland
Icepharma hf.
Tel: +354-540 8000

Slovak Republic
Bayer, spol. s r.o.
Tel: +421 2-59 21 31 11

Italy
Bayer S.p.A.
Tel: +39 02 397 81

Finland
Bayer Oy
Tel: +358-20 785 21

Cyprus
NOVAGEM Limited
Τηλ: +357 22 48 38 58

Sweden
Bayer AB
Tel: +46 (0)8 580 223 00

Latvija
SIA Bayer
Tel: +371 67 845 563

United Kingdom (Northern Ireland)
Bayer AG
Tel: +44-(0)118 206 3000

This leaflet was last approved in

Other sources of information

More information about this medicine is available on the website of the European Medicines Agency (<http://www.ema.europa.eu>).

Transtoyou