

Package leaflet: information for the user Mebendazole CF 100 mg, tablets mebendazole

Read carefully the leaflet before you take this medicine use, because there is important information in for you.

Use this medicine always exactly as described in this leaflet or as your doctor or pharmacist you that has told.

Keep this leaflet. Maybe you need it later again needed.

Do you still questions? Then contact your pharmacist.

Do you experience one of the side effects listed in section 4 appear? Or do you a side effect that not in this leaflet is? Then contact your doctor or pharmacist.

Does your complaint after 14 days not less, or does it even worse? Then contact your doctor.

Contents of this leaflet

1. What is Mebendazole CF 100 mg and what is this medicine usedfor
2. When may you this medicine not take or must you be extra careful with be?
3. How do you use this medicine ?Possible
4. side effects side effects
5. How do you store this medicine ?Contents
6. of the package and other information What is

1. Mebendazole CF 100 mg and what is it used for is this medicine drug used?

Therapeutic group

Mebendazole CF 100 mg belongs to the group of medicines that anthelmintics is called. Mebendazole is a drug that worms can kill. It is effective against infections of pinworms, roundworms, whip- and hookworms.

Mebendazole kills both worms, larvae as eggs.

In some cases (mainly severe infections with hookworms or whipworms) where no complete recovery obtained is, occurs there indeed a reduction of the egg excretion takes place of at least 90%.

Application of the medication

Mebendazole CF 100 mg is used for infections by pinworms, roundworms, whipworms and hookworms.

2. When should you not take this medicine of must you extra careful with be?

When may you this medicine not use?

- You are allergic to one of the ingredients in this medicine. These substances can you find in section 6.

When should you be extra careful with this medicine drug?

Contact your doctor or pharmacist before you use this medicine:

- In children younger than 2 years. There is little experience with the drug in children younger than 2 years; the use of mebendazole in children younger than 2 years is therefore not recommended for advise.
- There is no special diet or laxative to be used when applying mebendazole .. the worm infection worm infection accompanied goes with diarrhea, must one also that diarrhea treat.
- To reinfection after treatment to prevent, should in infection with pinworms the whole family to be treated.
- The excipient sunset yellow (E110) may cause allergic reactions , including, cause. Hypersensitivity reactions occur more frequently in individuals with hypersensitivity to acetylsalicylic acid acetylsalicylic acid.
- also use metronidazole metronidazole Combination. Combination of these products can lead to Stevens-Johnson syndrome (severe hypersensitivity reaction with (high) fever, red spots on the skin, joint pains eye infection/of eye inflammation) of toxic epidermal necrolysis (severe, sudden (hypersensitivity) reaction accompanied with fever and and blisters on the skin/peeling of the skin).

Do you use any other medications medicines Do you use any besides Mebendazole CF 100 mg any other other medicines, have you that recently done or is there a possibility that you in the near future other medicines will use use? Tell that then your doctor or pharmacist.

Other medicines can the effect of mebendazole influence. This applies also for medications that you only occasionally and sometimes use, for medications that you recently used have used and for medications that you without prescription can buy.

The absorption of some medications depends on the amount of acid in the stomach; mebendazole can the absorption of these medications affect.

In prolonged use of high doses can there be an interaction occur with the antacid cimetidine.

Pregnancy and breastfeeding

Are you pregnant, do you think you are pregnant to be, do you want to become pregnant or are you breastfeeding ? Then contact your? Then contact your doctor, before you this medicine use. Mebendazole can be used during the pregnancy, however during the first months only on advice of your doctor. The possible risks in the prescribing of Mebendazole CF 100 mg during the pregnancy should be weighed against become against the expected benefits of the treatment. During the period of breastfeeding only use after consultation with your doctor.

Ask your doctor or pharmacist for advice before you take a medicine ..

and the use of machines There

are no data data known about the effect of this product on the driving ability and the reaction time. An effect is however not likely.

Mebendazole CF 100 mg contains lactose

Has your doctor told that you some sugars do not tolerate? Then contact your doctor before you take this medicine.

Mebendazole CF 100 mg contains the colorant sunset yellow (E110)
Sunset Yellow can allergic reactions cause.

Mebendazole CF 100 mg contains sodium
This medicine contains less than 1 mmol sodium (23 mg) per tablet, which means that that it in essence 'sodium-free' is.

3. How do you use this medicine ?Use

this medication always exactly as described in in this leaflet or as your doctor or pharmacist you that has told. Are you unsure about the correct use? Then contact your doctor or pharmacist.

Dosage

The following dosage applies both for children from 2 years as for adults. There is no special diet of laxative to be prescribed. If the worm infection is accompanied by with diarrhea, one should simultaneously the symptoms of this of this to treat.

Infections with pinworms: 1 tablet and after 14 days again 1 tablet, if necessary every 14 days repeat. Infections with roundworms, whipworms and hookworms: 1 tablet in the morning and 1 tablet in the evening, for 3 consecutive days.

If after three weeks no complete cure is achieved, should the treatment be repeated ..

which way to take ?The

tablets tablets can with some water be taken or chewed.

Crush the tablet before you give it to a young child gives. Keep a child always in the holes while the this medicine takes.

Children younger than 2 years

Due to insufficient data no dosage advice can be given for children younger than 2 years. Do not give mebendazole to children under the 2 years without first consulting your doctor or pharmacist first tea have consulted.

Consult a doctor if the symptoms worsen or within 14 days do not improve.

In case you notices that mebendazole too strongly or just too little works, consult then your doctor or pharmacist.

Do you too much of this medicine used?

When you too much Mebendazole CF 100 mg, tablets have taken, contact immediately your doctor or doctor or pharmacist.

At overdose occur gastrointestinal complaints such as abdominal pain, nausea, vomiting and diarrhea occur. The patient water or lemonade let drink with therein activated charcoal. If the tablets shortly beforehand taken are, it is useful the patient to induce vomiting.

Did you forget this medicine to use?

In pinworms it is of great importance that the second dose after 14 days is taken. If this period longer than 15 days is must you probably still a third tablet take after 14 days. Consult in that case your doctor.

In the three-day courses take you the missed dose as soon as you think of it . However,when it is time time is for the next dose, take your usual dose at and complete the course until the tablets are finished are.

Take no double dose to make up for a missed dose in catching up.

Do you still have questions about the of this medication? Then contact your doctor or pharmacist.

4. Possible side effects

Like any medication can also this medication have side effects , althoughnot everyone experiences not everyone with it to make.

Below are the side effects listed that may occur with the treatment with Mebendazole CF 100 mg. The evaluation of the side effects is based on the following frequencies:

Very common: In more than 1 in 10 users Common: In 1 to 10 in 100 users

Sometimes: In 1 to 10 of the 1,000 users

Rarely: In 1 to 10 of the 10,000 users

Very rarely: In less than 1 of the 10,000 users

Not known: Can with the available data not be determined

General disorders

Not known: fever

Blood- and lymphatic system disorders

Rare: deficiency of white blood cells

Immune system disorders

Rarely: hypersensitivity, in severe hypersensitivity to substances in the medication can a severe reaction occur in which the blood vessels suddenly very wide become, with as result that the blood pressure drops and the heart rapidly but weakly beats, visible if pallor, restlessness, and clammy skin (anaphylactic reaction). In extreme cases can loss of consciousness occur (anaphylactoid reaction).

Nervous system disorders

Rare: convulsions, dizziness

Gastrointestinal disorders

Common: abdominal pain

Sometimes: stomach- and intestinal complaints, diarrhea, flatulence

Liver- and gallbladder diseases

Rarely: inflammation of the liver, abnormal liver function tests

Skin- and subcutaneous disorders

Rarely: itching, a skin condition where the outer layer of the skin detaches of the underlying layers and erythema multiforme, a condition where swollen red spots blisters can form (Stevens-Johnson syndrome or toxic epidermal necrolysis), skin rash, swollen face (angio-edema), hives, hair loss

Consult immediately your doctor if one of the following symptoms occur:

- A severe skin disorder, such as rash, blisters on the skin and painful sores in the mouth, eyes or ano-genital region (concerns the anus and genitals), accompanied by fever.
- A reaction that occurs immediately after administration and characterized is by skin rash, itching, shortness of breath and swollen face.
- A severe hypersensitivity reaction that occurs short after the administration and characterized is by hives, itching, flushing, fainting and breathing difficulties, in the midst of other symptoms.

The reporting of side effects

Do you experience side effects, please contact your doctor or pharmacist. This also applies to possible side effects side effects that not in this leaflet are listed. You can side effects also directly report via the Dutch Side Effects Centre Lareb, website: www.lareb.nl. By reporting side effects ,, can you us help more information to obtain about the safety of this medication.

5. How do you store this medication ??

Keep out of the sight and reach of children .. below 25 °C. Store

in the original packaging to protect against light and and moisture.

Use this medicine not anymore after the expiry date. It is to find on the packaging after "Do not to use after:" or "Exp.:". There is a month and a year . The last day of that month is the expiry date expiration date.

medicines medicines not through the sink or the toilet and throw them not in the trash bin. Ask your pharmacist what to do with medicines you should do not more used. They are then in a responsible manner destroyed and do not enter the environment end up.

6. Contents of the packaging and other information Which substances are in this medicine ?

The

- active ingredient in this medicine is mebendazole . Each tablet contains 100 mg mebendazole mebendazole.
- The other substances in this medicine are lactose monohydrate (see under section 2 "Mebendazole CF 100 mg contains lactose"), potato starch, sodium starch glycolate, saccharin-sodium, povidone (E1202), talc, colloidal anhydrous silicon dioxide (E551), magnesium stearate (E470b), orange flavoring, sunset yellow FCF aluminum lake (E 110) (see below section 2 "Mebendazole CF 100 mg contains the colorant sunset yellow FCF Aluminum lake (E110)").

How does Mebendazole CF 100 mg look and how much is in a packaging?

The tablets are pink-yellow to orange-yellow, round with a diameter of 10 mm with a score line on one side and on the other side optional mentioned "mebendazole" and "100".

Tablets in blister packaging. The blister strip contains 2, 6 or 10 tablets, packaged in multiples of 2, 6 or 10 tablets in a box.

Not all mentioned pack sizes are marketed in the market.

Holder of the authorization for the in the trade bringing and manufacturer

Marketing Authorization Holder

Centrafarm B.V.

Van de Reijtstraat 31-E

4814 NE Breda

Netherlands

Manufacturer

Centrafarm Services B.V.

Van de Reijtstraat 31-E

4814 NE Breda

Netherlands

Sanico NV

Veedijk 59

2300 Turnhout

Belgium

In the register registered under number

RVG 57042, Mebendazole CF 100 mg, tablets

This leaflet is for the last approved in December 2024

More information about this medication is available on the website of the CBG (www.cbg-meb.nl).